

How to Inject Andembry®

(garadacimab-gxii) — Self-Injection Guide for HAE Prevention



SCAN FOR FULL GUIDE

THE IMMUNE CONNECTION

Hereditary angioedema (HAE) attacks are driven by uncontrolled activation of the contact (kallikrein-kinin) pathway, which leads to excess bradykinin and swelling. Andembry works by blocking Factor XIIa, the enzyme that starts this cascade — helping prevent attacks before they begin.

WHAT IS ANDEMBRY?

Andembry is a prescription medicine given as a shot under the skin (subcutaneous injection) to help prevent attacks of hereditary angioedema (HAE). It comes as a prefilled autoinjector or a prefilled syringe with a needle safety device.

YOUR DOSING SCHEDULE

First dose (loading dose): 400 mg

Given as two separate 200 mg injections, on the same day, at two different sites.

Monthly maintenance dose: 200 mg

One injection, once a month, on the same date each month.

If you miss a dose:

Give the injection as soon as you remember, then continue your regular monthly schedule.

BEFORE YOU INJECT

- 1 Remove the autoinjector or syringe from the refrigerator and let it sit at room temperature for 30 minutes. Do not heat it or microwave it.
- 2 Do not shake the device.
- 3 Look at the liquid through the viewing window. It should be slightly opalescent to clear and brownish-yellow to yellow. Do not use if cloudy, discolored, or has particles.
- 4 Check the expiration date on the label. Do not use if expired.
- 5 Gather your supplies: alcohol wipe, cotton ball or gauze, and a sharps disposal container.

AFTER THE INJECTION

- Place the used autoinjector or syringe in a sharps disposal container right away.
- Do not throw the used device in the household trash.
- Do not reuse the autoinjector or syringe.
- Do not recap the needle.

WHY VEROS HEALTH?

- ✓ Hands-on self-injection training from our nursing team before you start at home.
- ✓ Direct access to your care team for questions on technique, side effects, or scheduling.
- ✓ Whole-patient HAE management, coordinated across our multi-specialty team.
- ✓ 30+ years of immune expertise — IMMUNOe Health & Research.

CHOOSING AN INJECTION SITE

Thigh (front):

A good option for self-injection.

Abdomen (belly):

Another good option. Stay at least 1 inch (2 cm) away from your belly button.

Upper arm (back):

Use only if a caregiver is giving the injection.

- Rotate your injection site each time you inject.
- Do not inject into skin that is tender, bruised, red, hard, or scarred.
- For the loading dose, use a different site for each of the two injections.

HOW TO GIVE THE INJECTION

- 1 Wash your hands thoroughly with soap and water.
- 2 Clean the injection site with an alcohol wipe and let it air dry.

Using the prefilled autoinjector:

- Remove the cap.
- Place it flat against your skin at the chosen site.
- Press and hold it firmly against your skin. You may hear a click as the injection starts.
- Keep holding it in place until the injection is complete (follow your device's timing).

Using the prefilled syringe with needle safety device:

- Remove the needle cap.
- Gently pinch a fold of skin at the injection site.
- Insert the needle at the angle shown in your device instructions.
- Slowly push the plunger all the way down to give the full dose.
- Release the skin fold and remove the needle.

If there is a small amount of bleeding, gently press a cotton ball or gauze over the site. Do not rub the area.

STORAGE

- Store Andembry in the refrigerator at 36°F to 46°F (2°C to 8°C).
- Keep it in the original carton to protect it from light.
- Do not freeze. Do not shake.

When to call your Veros provider: Call right away if you notice signs of an allergic reaction — rash, hives, swelling of the face or throat, or difficulty breathing. Also call with questions about your injection technique or dosing schedule, or if any side effect concerns you.